

Click-iT® RNA Imaging Kits

B066

Maiwen Caudron-Herger

m.caudron@dkfz-heidelberg.de

Click-iT® RNA Imaging Kits

Catalog nos. C10329, C10330

Table 1. Contents and storage information.

Material	C10329	C10330	Concentration	Storage*	Stability
5-ethynyl uridine (EU) (Component A)	10 mg	10 mg	NA	<ul style="list-style-type: none"> • 2–6°C • Protect from light • Dessicate • DO NOT FREEZE 	When stored as directed, this kit is stable for at least 1 year.
Alexa Fluor® azide (Component B)	1 vial (Alexa Fluor® 488 azide)	1 vial (Alexa Fluor® 594 azide)			
Click-iT® RNA reaction buffer (Component C)	25 mL	25 mL	1X		
Copper (II) sulfate (CuSO ₄) (Component D)	1 mL	1 mL	100 mM aqueous solution		
Click-iT® reaction buffer additive (Component E)	400 mg	400 mg	NA		
Click-iT® reaction rinse buffer (Component F)	50 mL	50 mL	1X		
Hoechst 33342 (Component G)	35 µL	35 µL	10 mg/mL in water		

*These storage conditions are appropriate when storing the entire kit upon receipt. After preparing stock solutions, optimal storage conditions may change. For storing prepared stock solutions, follow recommendations included in this product information sheet. NA = Not applicable.

Number of assays: Sufficient material is supplied for 25 coverslips, based on the protocol below.

Approximate fluorescence excitation/emission maxima: Alexa Fluor® 488 azide: 495/519 nm; Alexa Fluor® 594 azide: 590/615 nm; Hoechst 33342: 350/461 nm, bound to DNA.

Introduction

The Click-iT® RNA Imaging Assay enables detection of global RNA synthesis temporally and spatially in cells and tissues.¹ The ability to detect newly synthesized RNA or changes in RNA levels resulting from disease, environmental damage, or drug treatments is an important aspect of toxicological profiling. Utilizing an alkyne-modified nucleoside, 5-ethynyl uridine (EU), and powerful click chemistry, newly synthesized RNA can be detected without the use of radioactivity or antibodies with a simple, two-step procedure. In step one, the alkyne-containing nucleoside is fed to cells or animals, and is actively incorporated into nascent

RNA. The small size of the tag enables efficient incorporation of the modified nucleoside into RNA, but not into DNA. Detection utilizes the chemoselective ligation or “click” reaction between an azide and an alkyne where the modified RNA is detected with a corresponding azide-containing dye (Figure 1). With its diminutive “footprint”, the Click-iT® detection molecule can easily penetrate complex samples and leaves open the possibility of multiplex analyses with other probes, including antibodies for the detection of RNA-interactive proteins for deeper biological insights.

Click reactions have several general characteristics: the reaction is efficient, no extreme temperatures or solvents are required, the reaction is complete within 30 minutes, the components of the reaction are bioinert, and perhaps most importantly, no side reactions occur — the label and detection tags react selectively and specifically with one another.^{2–5} This final point is a key advantage of this powerful detection technique; it is possible to apply click chemistry-labeled molecules to complex biological samples and detect them with unprecedented sensitivity due to extremely low background.

The kits contain all of the components needed to label and detect newly synthesized RNA in whole cells (Figures 2–3). The kits are also supplied with blue fluorescent Hoechst 33342 dye as a nuclear counterstain or for DNA profiling. The kits include sufficient reagents for labeling 25, 18 × 18 mm coverslips using 500 µL reaction volume per well. For larger quantities, inquire at www.invitrogen.com.




Figure 1. Click reaction between EU and azide-modified dye or hapten..




Figure 2. Workflow diagram for the Click-iT® RNA Imaging Assay.




Figure 3. Dose response for actinomycin D in HeLa cells using the Click-iT® RNA Assay. HeLa cells were treated with the indicated amounts of actinomycin D for 18 hours, followed by a 1 hour incubation with 5-ethynyl uridine (EU). Cells were then fixed and permeabilized, and EU incorporated into newly synthesized RNA was detected using the green-fluorescent Alexa Fluor® 488 azide. Quantitative analysis was performed using the Thermo Scientific Cellomics® ArrayScan® VTI and Compartmental Analysis Bioapplication.

Table 2. Click-iT® detection reagent compatibility.

Molecule	Compatibility*
Qdot® nanocrystals	Use Qdot® nanocrystals after the Click-iT® detection reaction.
Fluorescent proteins such as Green Fluorescent Protein (GFP)	Use organic dye-based reagents, such as TC-FIAsH™ or TC-ReAsH™ reagents, for protein expression detection or anti-GFP rabbit or chicken antibodies before the Click-iT® detection reaction.
Organic dyes such as Alexa Fluor® dyes, fluorescein (FITC)	Completely compatible with the Click-iT® detection reaction.
TC-FIAsH™ or TC-ReAsH™ reagents	Detect the tetracysteine (TC) tag with FIAsH™ or ReAsH™ reagents before the Click-iT® detection reaction.
Phalloidin	Phalloidin staining is not compatible with the Click-iT® detection reaction. Use antibodies against other proteins, such as anti- α -tubulin, for visualization of the cytoskeleton.
Horseradish peroxidase (HRP)	Use HRP after the Click-iT® detection reaction.
R-phycoerythrin (R-PE) and R-PE-tandems such as Alexa Fluor® 680-R-PE	Use R-PE and R-PE tandems after the Click-iT® detection reaction.
Allophycocyanin (APC) and APC-tandems	Completely compatible with the Click-iT® detection reaction.

*Compatibility indicates whether the fluorescent molecule itself or the detection method involves components that are unstable in the presence of copper catalyst used for the Click-iT® detection reaction.

Before You Begin

Materials Required but Not Provided

- Phosphate buffered saline (PBS), pH 7.2–7.6
- Fixative (*i.e.*, 3.7% Formaldehyde in PBS)
- Permeabilization reagent (*i.e.*, 0.1% Triton® X-100 in PBS)
- Deionized water (dH₂O)
- Dimethylsulfoxide (DMSO)
- *Optional*: 6-well plates

Cautions

Hoechst 33342 (Component G) is a known mutagen. Use the dye with appropriate precautions.

DMSO is known to facilitate the entry of organic molecules into tissues. Handle reagents containing DMSO using equipment and practices appropriate for the hazards posed by such materials. Dispose of the reagents in compliance with all pertaining local regulations

Preparing the Stock Solutions

- 1.1 To prepare a 100 mM stock solution of EU (Component A), add 373 μ L of deionized water to the vial and mix until fully dissolved.

After use, store any remaining stock solution at $\leq -20^{\circ}\text{C}$. When stored as directed, this stock solution is stable for up to 1 month.

Note: Additional EU is available separately from Invitrogen (Cat. no. E10345).

1.2 To prepare a stock solution of the Alexa Fluor® azide:

- Add 94 µL of DMSO to Alexa Fluor® 488 vial (Component B, Cat. no. C10329), and mix well by pipetting or vortexing.
- Add 85 µL of DMSO to Alexa Fluor® 594 vial (Component B, Cat. no. C10330), and mix well by pipetting or vortexing.

After use, store any remaining stock solution at $\leq -20^{\circ}\text{C}$. When stored as directed, this stock solution is stable for up to 1 year.

1.3 To prepare a 10X stock solution of the Click-iT® reaction buffer additive (Component E), add 2 mL of deionized water to the vial and mix until fully dissolved.

After use, store any remaining stock solution at $\leq -20^{\circ}\text{C}$. When stored as directed, this stock solution is stable for up to 1 year. If the solution develops a brown color, it has degraded and should be discarded.

Experimental Protocols

Labeling Cells with EU

The following protocol was developed with HeLa and NIH3T3 cells, plated at 40–50% confluency, and fed the following day with 1 mM EU for 1 hour. Growth medium, cell density, cell type variations, and other factors may influence the assay conditions. In initial experiments, we recommend testing a range of EU concentrations to determine the optimal concentration for your cell type and experimental conditions. Sufficient material is supplied for 25 coverslips based upon the protocol below.

- 2.1 Plate cells at desired confluency and allow cells to recover before additional treatment.
- 2.2 *Optional:* Perform cell treatments as desired. To address the potential reversibility of drug action on RNA synthesis, we recommend that you do **not** remove the drug-containing media during EU treatment.
- 2.3 Prepare a 2X working solution of EU from the 100 mM stock solution (prepared in step 1.1) in **pre-warmed** complete medium. For example, for a 1 mM final EU treatment, prepare a 2 mM working solution. Add an equal volume of this 2X EU working solution to the media containing cells.
- 2.4 Incubate under normal cell culture conditions for 1 hour. The optimal incubation time depends on the cell growth rate and cell treatment, and needs to be determined experimentally.
- 2.5 Proceed immediately to **Cell Fixation, Permeabilization, and Click-iT® Detection**.

Cell Fixation, Permeabilization, and Click-iT® Detection

The following protocol is based upon volumes convenient for processing the coverslips in a 6-well plate, with each well containing a single coverslip, but may be modified to fit other experimental formats. It is optimized with a fixation step using 3.7% formaldehyde in PBS followed by a 0.5% Triton® X-100 permeabilization step, but is amenable to other fixation and permeabilization reagents such as methanol and saponin.

- 3.1 Transfer the coverslips into a 6-well plate.
- 3.2 Add 1 mL of 3.7% formaldehyde in PBS to each well and incubate for 15 minutes at room temperature.

- 3.3** Remove the fixative and wash each well once with PBS.
- 3.4** Remove the wash solution and add 1 mL of 0.5% Triton® X-100 in PBS to each well. Incubate for 15 minutes at room temperature.
- 3.5** Prepare a 1X working solution of Click-iT® reaction buffer additive by diluting the 10X solution (prepared in step 1.3, see Table 3 for volumes) 1:10 in deionized water. Prepare this solution **fresh** and use the solution on the same day.
- 3.6** Prepare Click-iT® reaction cocktail according to Table 3.

Note: Add the components of the Click-iT® reaction cocktail in the order listed in Table 3, and use the cocktail immediately after preparation. The Click-iT® reaction buffer additive is susceptible to oxidation and is the limiting factor to the Click-iT® reaction cocktail's effectiveness over time.

Table 3. Click-iT® reaction cocktails.

Reaction components	Number of coverslips					
	1	2	4	5	10	25
Click-iT® RNA reaction buffer (Component C)	428 µL	856 µL	1.7 mL	2.1 mL	4.3 mL	10.7 mL
CuSO ₄ (Component D)	20 µL	40 µL	80 µL	100 µL	200 µL	500 µL
Alexa Fluor® azide (prepared in step 1.2)	1.8 µL	3.7 µL	7.4 µL	9.3 µL	18.8 µL	47 µL
Click-iT® reaction buffer additive (prepared in step 3.5)	50 µL	100 µL	200 µL	250 µL	500 µL	1.25 mL
Approximate total volume	500 µL	1 mL	2 mL	2.5 mL	5 mL	12.5 mL

- 3.7** Remove the permeabilization buffer (step 3.4) and wash each well once with PBS.
- 3.8** Remove the wash solution and add 500 µL of Click-iT® reaction cocktail (prepared in step 3.6) to each sample. Incubate for 30 minutes at room temperature, **protected from light**.
- 3.9** Remove the Click-iT® reaction cocktail and wash cells once with 1 mL of Click-iT® reaction rinse buffer (Component F). Then remove the Click-iT® reaction rinse buffer.
- 3.10** *Optional:* Perform antibody labeling on the samples at this time, following the recommendations from the manufacturer of the primary and secondary antibody. It is important to keep the samples **protected from light** during incubations.

For DNA staining and cell cycle distribution, proceed to **DNA Staining**. If no additional staining is desired, proceed to **Imaging and Analysis**.

DNA Staining The following protocol is based upon 1 mL of Hoechst 33342 working solution per sample.

- 4.1** Wash samples with PBS, then remove the wash solution.
- 4.2** Dilute Hoechst 33342 (Component G) 1:1,000 in PBS.
- 4.3** Add 1 mL of diluted Hoechst 33342 to each well, and incubate for 15 minutes at room temperature, **protected from light**.
- 4.4** Remove the Hoechst 33342 solution, and wash cells twice with PBS.
- 4.5** Remove the wash solution and proceed to **Imaging and Analysis**.

Imaging and Analysis

Click-iT® EU stained cells are compatible with all methods of slide preparation including wet mount or prepared mounting media. Image and analyze using appropriate instrument filter sets.

See Table 4 for the appropriate fluorescence excitation/emission maxima for Alexa Fluor® dyes and Hoechst 33342 bound to DNA.

Table 4. Approximate fluorescence excitation/emission maxima.

Fluorophore	Excitation (nm)	Emission (nm)
Alexa Fluor® 488	495	519
Alexa Fluor® 594	590	615
Hoechst 33342, bound to DNA	350	461

References

1. Proc Natl Acad Sci USA 105, 15779 (2008); 2. Chem Bio Chem 4, 1147 (2003); 3. J Am Chem Soc 124, 3192 (2003); 4. Angew Chem Int Ed Engl 41, 2596 (2002); 5. Angew Chem Int Ed Engl 40, 2004 (2001).

Product List Current prices may be obtained from our website or from our Customer Service Department.

Cat. no.	Product Name	Unit Size
C10329	Click-iT® RNA Alexa Fluor® 488 Imaging Kit *for 25 coverslips*.....	1 kit
C10330	Click-iT® RNA Alexa Fluor® 594 Imaging Kit *for 25 coverslips*.....	1 kit
Related Products		
C10289	Click-iT® AHA Alexa Fluor® 488 Protein Synthesis HCS Assay *2-plate size*.....	1 kit
C10327	Click-iT® RNA Alexa Fluor® 488 HCS Assay *2-plate size*.....	1 kit
C10328	Click-iT® RNA Alexa Fluor® 594 HCS Assay *2-plate size*.....	1 kit
C10337	Click-iT® EdU Alexa Fluor® 488 Imaging Kit *for 50 coverslips*.....	1 kit
C10339	Click-iT® EdU Alexa Fluor® 594 Imaging Kit *for 50 coverslips*.....	1 kit
C10340	Click-iT® EdU Alexa Fluor® 647 Imaging Kit *for 50 coverslips*.....	1 kit
C10350	Click-iT® EdU Alexa Fluor® 488 HCS Assay *2-plate size*.....	1 kit
C10351	Click-iT® EdU Alexa Fluor® 488 HCS Assay *10-plate size*.....	1 kit
C10354	Click-iT® EdU Alexa Fluor® 594 HCS Assay *2-plate size*.....	1 kit
C10355	Click-iT® EdU Alexa Fluor® 594 HCS Assay *10-plate size*.....	1 kit
C10356	Click-iT® EdU Alexa Fluor® 647 HCS Assay *2-plate size*.....	1 kit
C10357	Click-iT® EdU Alexa Fluor® 647 HCS Assay *10-plate size*.....	1 kit
E10345	5-ethynyl uridine (EU)	5 mg
H3570	Hoechst 33342, trihydrochloride, trihydrate *10 mg/mL solution in water*	10 mL

Contact Information

Molecular Probes, Inc.

29851 Willow Creek Road
Eugene, OR 97402
Phone: (541) 465-8300
Fax: (541) 335-0504

Customer Service:

6:00 am to 4:30 pm (Pacific Time)
Phone: (541) 335-0338
Fax: (541) 335-0305
probesorder@invitrogen.com

Toll-Free Ordering for USA:

Order Phone: (800) 438-2209
Order Fax: (800) 438-0228

Technical Service:

8:00 am to 4:00 pm (Pacific Time)
Phone: (541) 335-0353
Toll-Free (800) 438-2209
Fax: (541) 335-0238
probestech@invitrogen.com

Invitrogen European Headquarters

Invitrogen, Ltd.
3 Fountain Drive
Inchinnan Business Park
Paisley PA4 9RF, UK
Phone: +44 (0) 141 814 6100
Fax: +44 (0) 141 814 6260
Email: euroinfo@invitrogen.com
Technical Services: eurotech@invitrogen.com

**For country-specific contact information,
visit www.invitrogen.com.**

Further information on Molecular Probes products, including product bibliographies, is available from your local distributor or directly from Molecular Probes. Customers in Europe, Africa and the Middle East should contact our office in Paisley, United Kingdom. All others should contact our Technical Service Department in Eugene, Oregon.

Molecular Probes products are high-quality reagents and materials intended for research purposes only. These products must be used by, or directly under the supervision of, a technically qualified individual experienced in handling potentially hazardous chemicals. Please read the Material Safety Data Sheet provided for each product; other regulatory considerations may apply.

Limited Use Label License No. 223: Labeling and Detection Technology

The manufacture, use, sale or import of this product may be subject to one or more patents or pending applications owned or licensed by Invitrogen Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity) in a manner consistent with the accompanying product literature. The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. The buyer may transfer information or materials made through the use of this product to a scientific collaborator, provided that such transfer is not for any Commercial Purpose, and that such collaborator agrees in writing (a) to not transfer such materials to any third party, and (b) to use such transferred materials and/or information solely for research and not for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For products that are subject to multiple limited use label licenses, the most restrictive terms apply. Invitrogen Corporation will not assert a claim against the buyer of infringement of patents that are owned or controlled by Invitrogen Corporation and/or Molecular Probes, Inc. which cover this product based upon the manufacture, use or sale of a therapeutic, clinical diagnostic, vaccine or prophylactic product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. If the purchaser is not willing to accept the limitations of this limited use statement, Invitrogen is willing to accept return of the product with a full refund. For information on purchasing a license to this product for purposes other than research, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Several Molecular Probes products and product applications are covered by U.S. and foreign patents and patents pending. All names containing the designation ® are registered with the U.S. Patent and Trademark Office.

Copyright 2009, Molecular Probes, Inc. All rights reserved. This information is subject to change without notice.