

Blood Clotting and Bowel Cancer Risk

People whose blood clots more easily than normal are more often affected by cancer. Scientists at the German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ) have discovered that a number of variants of clotting factor genes have an influence on bowel cancer risk. They found out that carriers of a particular gene variant of clotting factor V have a bowel cancer risk that is six times higher than people who do not have this gene variant. This study is an important basis for finding out whether and for whom drugs which affect blood clotting may be useful to prevent bowel cancer.

Back in the mid 19th century, a French doctor, Armand Trousseau, discovered a connection between cancer and thrombosis – the formation of often dangerous blood clots that can lead to venous occlusion. Today it is known that cancer and its treatment change blood flow properties and thus promote the formation of clots. However, clots do not only occur as a side effect and consequence of cancer, but, vice versa, an increased blood clotting tendency may also be associated with an elevated cancer risk.

About twelve different blood proteins called clotting factors interact in a coordinated manner in the blood clotting (coagulation) process. In the same way as hemophilia (decreased blood clotting) is inherited, genes also play a role in an increased clotting tendency (thrombophilia): There are well studied gene variants (polymorphisms) of a number of clotting factors which are associated with an increased or decreased clotting tendency. Between two and five percent of the population carry such genetic variants.

At the German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ) in Heidelberg, scientists headed by Professor Dr. Hermann Brenner have been studying six gene variants of different clotting factors for a possible connection with colorectal cancer risk. In a large study, they analyzed the occurrence of these six variants in approximately 1.800 colorectal cancer patients and in the same number of healthy control persons.

The team found the most obvious connection for a variant that substantially increases the risk of thrombosis and which is known as factor V Leiden (FVL). Study participants who carry this genetic variant on both copies of their chromosome 1 were found to have a six fold increase in colorectal cancer risk compared to participants who carry two copies of the “standard variant” of factor V. If only one copy of chromosome 1 had the FVL variant, bowel cancer risk was not elevated.

Another connection with bowel cancer prevalence was found by the research team for a particular gene variant of clotting factor XIII: People with this mutation are slightly more rarely affected by venous thrombosis than those who carry the factor XIII standard version. Now the DKFZ team has shown that their colorectal cancer risk is also 15 percent lower. For the other four gene variants studied the team found no connection with bowel cancer risk.

It is known today that coagulation and carcinogenesis are associated. Thus, the interplay of all coagulation (clotting) factors leads to the formation of active thrombin, which, in turn, activates hemostatic fibrin. However, thrombin also contributes to the formation of new blood vessels and is able to dissolve the extracellular matrix, which is the adhesive that keeps cells together. Thrombin may thus make it easier for cancer cells to invade surrounding tissue.

“It is interesting that not every gene variant that increases the tendency to clot automatically also increases the risk of developing colorectal cancer. It also makes a difference whether the gene variant is present on both chromosomes or just on one of them. We therefore have to analyze in detail which clotting factors affect cancer risk and in what manner,” explained study head Hermann Brenner. The knowledge of these connections is the first prerequisite for finding out whether and for whom drugs that affect blood clotting may prevent bowel cancer.

Carla Y. Vossen, Michael Hoffmeister, Jenny C. Chang-Claude, Frits R. Rosendaal, and Hermann Brenner: Clotting Factor Gene Polymorphisms and Colorectal Cancer Risk. *Journal of Clinical Oncology* 2011, DOI: 10.1200/JCO.2010.31.8873

The German Cancer Research Center (Deutsches Krebsforschungszentrum, DKFZ), employing over 2,500 staff members, is the largest biomedical research institute in Germany. More than 1,000 scientists are working to investigate the mechanisms of cancer development, identify cancer risk factors and develop new strategies for better cancer prevention, more precise diagnosis and effective treatment of cancer patients. In addition, the staff of the Cancer Information Service (KID) provides information about this widespread disease for patients, their families, and the general public. DKFZ is funded by the German Federal Ministry of Education and Research (90%) and the State of Baden-Wuerttemberg (10%) and is a member of the Helmholtz Association of National Research Centers.

Dr. Stefanie Seltsmann
Head of Press and Public Relations
German Cancer Research Center
Im Neuenheimer Feld 280
D-69120 Heidelberg
T: +49 6221 42 2854
F: +49 6221 42 2968
presse@dkfz.de